



Hampton Roads Gastroenterology

303 Butler Farm Road Suite 103, Hampton VA 23666

Tel: (757) 826-3434 Fax: (757) 826-9028

MoviPrep Bowel Prep for Colonoscopy

Please read all prep instructions, as well as the Procedure Instructions also provided.

- Obtain your prep kit from the pharmacy as soon as possible.
- If you have any problems with your prep, please call the office at 826-3434.
- **DO NOT FOLLOW THE INSTRUCTIONS PROVIDED BY THE PACKAGE INSERT; FOLLOW THESE INSTRUCTIONS FOR YOUR PREP.** Thank you!

For the 3 days prior to your procedure: Avoid greens, vegetables, leafy foods, corn, seeds, & fruits with seeds

On the day before the procedure:

1. You may only have CLEAR LIQUIDS. Try to drink something every hour that you are awake.
You may have: apple juice, white grape juice, Jell-O (ONLY green or yellow), popsicles (ONLY green or yellow), 7-UP, Sprite, Ginger Ale, weak tea, CLEAR broths (with no food in it), and bouillon cubes.
You MAY NOT have: orange juice, coffee, Coke, Pepsi, solid foods, milk, any broths with bits of food or highly seasoned broths, or alcoholic beverages.
2. If you are *extremely* hungry, you may have a cup of blended yogurt (no fiber) or a nutrition drink such as Boost or Ensure before 2 PM.
3. Between 5 and 7 PM, you will begin the MoviPrep. Empty 1 pouch A and 1 pouch B into the disposable container. Add lukewarm drinking water to the top line of the container. Mix to dissolve. This can be done 24 hours in advance and refrigerated).
4. The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz.), until the full liter is complete.
5. Drink 16 ounces of the clear liquid of your choice.

On the morning of your procedure:

1. Three hours before your scheduled procedure time, but no later than 7 AM, empty 1 pouch A and 1 pouch B into the disposable container. Add lukewarm drinking water to the top line of the container. Mix to dissolve. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz.), until the full liter is complete. Drink 16 ounces of the clear liquid of your choice.
2. **STOP DRINKING ALL LIQUIDS 3 HOURS BEFORE YOUR SCHEDULED PROCEDURE TIME. YOU MAY NOT EAT ANY FOOD ON THE DAY OF YOUR PROCEDURE.**

You must have a driver for your procedure; you will be sedated and not allowed to drive yourself. You may not take a taxi or bus home from the procedure. If you arrive without a driver, you may be charged a \$150 fee.



Hampton Roads Gastroenterology

303 Butler Farm Road Suite 103, Hampton VA 23666

Tel: (757) 826-3434 Fax: (757) 826-9028

Procedure Instructions

! STOP ALL CLEAR LIQUIDS (including water) 3 HOURS BEFORE YOUR ARRIVAL TIME !

For endoscopies: NO FOOD THE DAY OF YOUR PROCEDURE

For colonoscopies: NO FOOD THE DAY BEFORE & DAY OF YOUR PROCEDURE

If you have any questions, please call the office at 757-826-3434.

- The day of your procedure, take medications as usual for blood pressure, asthma, seizures, anxiety, and other cardiac related drugs (i.e. beta blockers, calcium channel blockers). **If your procedure is early in the morning, take your pills with a small sip of water as soon as you get up.**
- Bring any prescribed inhalers with you to your procedure.
- If you are on a blood thinner such as **Coumadin, Warfarin, Eliquis, Plavix, Pradaxa, Xarelto, Effient, or Brilinta**, let your physician know before scheduling and they will determine whether to take it or not.
- If you are diabetic, we recommend that you take ½ of your usual insulin dose the day of your procedure. **Do not take oral medications for diabetes** (Metformin, Glucophage, etc.) **the morning of your procedure.** Please wait until after the procedure. We will check your blood sugar prior to the procedure. Optimal blood sugar for procedure is <250.
- Use of GLP-1 agonists for weight loss (*Ozempic semaglutide, Trulicity dulaglutide, Byetta exenatide, Victoza liraglutide, Bydureon exenatide, Wegovy semaglutide, Mounjaro/Zepbound tirzepatide, Rysbelsus semaglutide, Saxenda liraglutide*) must be stopped 1 week prior to the procedure.
- Use of phentermine must be stopped 3 days prior to the procedure.
- **Do not chew gum or candy the day of your procedure.**
- **DO NOT SMOKE MARIJUANA THE DAY OF YOUR PROCEDURE.**

You must have a driver for your procedure; you will be sedated and not allowed to drive yourself. You may not take a taxi, bus, or Uber/Lyft home from the procedure.

72 hours (3 business days) notice is required for cancellations or you will be charged a fee of \$150. We appreciate your cooperation.